CS 323: Numerical Analysis and Computing

Midterm #2

Copyright Notice: All course materials, including this exam, are copyrighted and owned by Prof. Mridul Aanjaneya. Posting/copying these materials to websites, such as Course Hero, is strictly prohibited. Violators will automatically fail the class.

Instructions: This is a **closed notes** exam. You are **not allowed** to consult any textbook, your class notes, homeworks, or any of the handouts from us. You are also **not permitted** to use laptop computers, cell phones, tablets, or any other hand-held electronic devices.

Name	
NetID	

Part #1	
Part #2	
Part #3	
Part #4	
Part #5	
TOTAL	

1. [24% = 4 questions × 6% each] MULTIPLE CHOICE SECTION. Circle or underline the correct answer (or answers). No justification is required for your answer(s).

   (a) Given a non-singular $n \times n$ system of linear equations $Ax = b$, the solution vector x remains unchanged after

   (Circle the TWO most correct answers.)

   i. Permuting the rows of A and b.
   ii. Permuting the columns of A.
   iii. Multiplying both sides of the equation from the left by a non-singular $n \times n$ matrix M.

   (b) Let A be an $m \times n$ matrix. Then the matrix $A^T A$ is always

   (Circle the TWO most correct answers.)

   i. symmetric.
   ii. non-singular.
   iii. positive definite.
   iv. positive semi-definite.

   (c) Suppose that an $n \times n$ matrix A is perfectly well-conditioned, i.e., the condition number $\kappa(A) = 1$ in the L_∞-norm. Which of the following matrices would then necessarily share this same property?

   (Circle the TWO most correct answers.)

   i. cA, where c is any non-zero scalar.
   ii. BA, where B is any non-singular matrix.
   iii. A^{-1}, the inverse of A.
   iv. DA, where D is a non-singular diagonal matrix.

   (d) What is the condition number of the following matrix using the L_1-norm?

   (Circle the ONE most correct answer.)

$$\begin{bmatrix} 4 & 0 & 0 \\ 0 & -6 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

   i. 2
   ii. 3
   iii. 1
   iv. 12

2. [18% = 3 questions × 6% each] SHORT ANSWER SECTION. Answer each of the following questions in no more than 2-3 sentences.

   (a) Suppose you have already solved the $n \times n$ linear system $Ax = b$ by LU factorization. What is the further cost (order of magnitude will suffice) of solving a new system

   - With the same matrix A, but a different right hand side vector b'?
   - With a different matrix A', but the same right hand side vector b?

   (b) Suppose the second column of the matrix $M_1 A$ during the LU decomposition algorithm is as follows:

$$a_2 = \begin{bmatrix} 3 \\ 2 \\ -1 \\ 4 \end{bmatrix}$$

   Specify the elimination matrix M_2 that zeros out the **last two** components of a_2.

   (c) If A and B are $n \times n$ matrices, with A non-singular and $c \in \mathbb{R}^n$, how would you efficiently compute the product $A^{-1}Bc$? (A need not be diagonally dominant.)

3. [16%] Consider the matrix

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 1 & 3 & 2 \end{bmatrix}$$

(a) Show that A is singular.

(b) If $b = (2, 4, 6)^T$, how many solutions are there to the system $Ax = b$?
Hint: For part (b), you just need to show if $b \in \mathsf{range}(A)$. You are not required to compute the entire solution set. So the answer should be either zero or infinite.

4. [12%] Let A be a rectangular $m \times n$ matrix with linearly independent columns, where $m > n$, and $b \in \mathbb{R}^m$. Show that for every $x \in \mathbb{R}^n$, we have

$$\|Ax - b\|_2^2 = \|A(x - x_0)\|_2^2 + \|Ax_0 - b\|_2^2$$

where $x_0 \in \mathbb{R}^n$ is the *least squares solution* of $Ax = b$.

Hint: Use the identity $\|x\|_2^2 = x^T x$.

5. [10%] Prove that the matrix

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

has no LU factorization, i.e., no lower triangular matrix L and upper triangular matrix U exist, such that $A = LU$.

Hint: Assume that there is an LU decomposition

$$\begin{bmatrix} l_{11} & 0 \\ l_{21} & l_{22} \end{bmatrix} \begin{bmatrix} u_{11} & u_{12} \\ 0 & u_{22} \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

and arrive at a contradiction!

6. [20%] Consider the $n \times n$ linear system $Ax = b$.

   (a) Show that if A is diagonal, the Jacobi method converges after just one iteration.

   (b) Show that if A is lower triangular, the Gauss-Seidel method converges after just one iteration.

   Hint: Decompose $A = D - L - U$ and use the definitions of the Jacobi and Gauss-Seidel methods, as described in the notes.